

CONIC-CONNECTED MANIFOLDS

PALTIN IONESCU* AND FRANCESCO RUSSO**

ABSTRACT. We study a particular class of rationally connected manifolds, $X \subset \mathbb{P}^N$, such that two general points $x, x' \in X$ may be joined by a conic contained in X . We prove that these manifolds are Fano, with $b_2 \leq 2$. Moreover, a precise classification is obtained for $b_2 = 2$. Complete intersections of high dimension with respect to their multi-degree provide examples for the case $b_2 = 1$. The proof of the classification result uses a general characterization of rationality, in terms of suitable covering families of rational curves.

A Gaetano Scorza che un secolo fa aveva colto l'importanza delle varietà razionalmente connesse considerando la classe particolare di varietà conicamente connesse formata da quelle di ultima specie.

"Invece per le V_4 di prima e terza specie arrivo a caratterizzarle tutte valendomi della teoria dei sistemi lineari sopra una varietà algebrica e, per le ultime, della circostanza che esse contengono un sistema ∞^6 di coniche così che per ogni loro coppia di punti passa una e una sola conica. Inoltre la natura dei ragionamenti è tale da mostrare come i risultati ottenuti possano estendersi, almeno per la maggior parte, alle varietà (di prima e ultima specie) a un numero qualunque di dimensioni, ...", [Sc2, Opere Scelte, vol. 1, p. 253].

INTRODUCTION

A projective manifold X is rationally connected if any two given (general) points of it may be joined by a rational curve. Since their introduction by Kollár, Miyaoka, and Mori [KMM], rationally connected manifolds were intensively studied, as they naturally generalize both classical notions of unirational and Fano manifold. When such an X is embedded in some projective space, a measure of its complexity is given by the minimal degree of a family of rational curves possessing the above defining property. As a first instance we have linear spaces, characterized by the presence of a line through any two distinct points. The next case is what we call *conic-connected* manifolds: They are embedded manifolds $X \subset \mathbb{P}^N$ such that for any two general points $x, x' \in X$, there is some irreducible conic joining x, x' and contained in X .

2000 *Mathematics Subject Classification.* 14M99, 14J45, 14E30.

Key words and phrases. Rationally connected, conic-connected, Fano manifold, quasi-line.

*Partially supported by the Italian Programme "Incentivazione alla mobilità di studiosi stranieri e italiani residenti all'estero".

**Partially supported by CNPq (Centro Nacional de Pesquisa), Grants 300961/2003-0, 308745/2006-0 and 474475/2006-9, and by PRONEX/FAPERJ–Algebra Comutativa e Geometria Algebrica.

The main results of this paper, Theorem 2.2 and Corollary 2.3, show that, over $\mathbb{C}$, conic-connected manifolds are Fano, with second Betti number $b_2 \leq 2$. Moreover, those with $b_2 = 2$ are completely described in the following short list: Segre products of two projective spaces and their hyperplane sections, or the inner projections, from a linear space, of the Veronese variety $v_2(\mathbb{P}^n)$. When $b_2 = 1$, unless $X \simeq v_2(\mathbb{P}^n)$, the Picard group is generated by the hyperplane section and the index of X is at least $\frac{\dim(X)+1}{2}$. Conversely, from a recent result by Bonavero–Höring, see [BH], it follows that any n -dimensional complete intersection which is Fano, of index at least $\frac{n+1}{2}$, is conic-connected; see Proposition 2.4. Moreover, the work of Hwang–Kebekus, see [HK], shows that any Fano n -fold of index greater than $\frac{2n}{3}$ is conic-connected; see Proposition 2.5. Applications of Theorem 2.2 to the classification of secant and dual defective manifolds, special Cremona transformations, etc. may be found in [Ru, IR].

The proof of Theorem 2.2 naturally fits into the context of Mori Theory, as established by the ground-breaking papers [Mo1, Mo2]. Besides the use of such powerful tools as the Cone Theorem, contractions of extremal rays, etc., we also need the very efficient method of algebraizing the equivalence relation associated to a covering family of curves. This technique, pioneered by Campana [Ca], was further developed and successfully applied by Kollár, Miyaoka, and Mori in their seminal paper [KMM]. The recent characterization of projective space, belonging to the same circle of ideas, and due to Cho, Miyaoka, and Shepherd-Barron (see [CMSB]), is also used in the proof of Theorem 2.2. Finally, a key ingredient of the argument is our Theorem 1.5, a “global version” of Theorem 1.3.

Theorem 1.3 which, we hope, has some independent interest, generalizes [IN, Theorem 4.2] (see also [Io2]). Its content is a criterion of rationality: X is rational if and only if it admits a covering family of rational 1-cycles, all passing through x , all smooth at x and such that the general cycle of the family is a curve, uniquely determined by its tangent line at x . In Theorem 1.5, under suitable assumptions, the birational isomorphism between X and $\mathbb{P}^n$ is shown to be biregular along a general (smooth) curve of the given family; moreover, the general member of the family corresponds to a line in $\mathbb{P}^n$.

We acknowledge our intellectual debt to the pioneering work of G. Scorza [Sc1, Sc2], as well as to all the previous papers mentioned above.

We work over $\mathbb{C}$; note however that Theorem 1.3 holds in any characteristic.

1. A RATIONALITY CRITERION

Let X be a projective variety. A family of rational 1-cycles on X , $\mathcal{F} \rightarrow \mathcal{C}$, is given by a closed subset $\mathcal{C} \subset \text{Chow}(X)$ such that for each 1-cycle of the family, all its irreducible components are rational. Moreover, we always assume that $\mathcal{C}$ is *irreducible* and the *general member* of $\mathcal{C}$ is an *irreducible reduced* (rational) curve.

Definition 1.1. A *covering family* (of rational 1-cycles, as above) on X is a family $\mathcal{F} \rightarrow \mathcal{C}$ as above, such that the tautological morphism $\mathcal{F} \rightarrow X$ is surjective.

Let $x \in X$ be a *smooth* point.

Definition 1.2. An x -covering family on X is a covering family $\mathcal{F} \rightarrow \mathcal{C}$ as above, such that, moreover, $x \in \text{Supp}(C)$ for any $C \in \mathcal{C}$ and the general member of $\mathcal{C}$ is smooth at x .

An x -covering family is called *maximal* if $\mathcal{C}$ is an irreducible component of the closed subscheme of $\text{Chow}_x(X)$, whose points correspond to rational 1-cycles.

An x -covering family is said to be *smooth* if all its 1-cycles are smooth at x .

An x -covering family is said to satisfy the *infinitesimal uniqueness property* if the general member of the family is uniquely determined by its tangent space at x .

Recall that X is *rationally connected* if through any two (general) points of X there passes a rational curve. Saying that X is rationally connected is equivalent to the existence of an x -covering family. One can also see that the existence of an x -covering family satisfying the infinitesimal uniqueness property is equivalent to X being *unirational*. The next result gives a criterion of *rationality*, in the spirit of the papers [IN, Io2]. It may be seen as a birational version of a series of characterizations of $\mathbb{P}^n$ in terms of families of rational curves, started in Mori's famous paper [Mo1].

Theorem 1.3. *Let X be a projective variety. The following conditions are equivalent:*

- (i) X is rational;
- (ii) for some $x \in X$, X admits a smooth x -covering family satisfying the infinitesimal uniqueness property.

Proof. To see that (i) implies (ii), note that a neighbourhood of a general point $x \in X$ is isomorphic to some neighbourhood of a point $y \in \mathbb{P}^n$. Simply consider the family induced on X by the lines of $\mathbb{P}^n$ passing through y .

(ii) implies (i), the non-trivial implication, is proved in the following three steps.

Step I. The basic diagram.

Let $\pi : \mathcal{F} \rightarrow \mathcal{C}$ be our x -covering family and let $\varphi : \mathcal{F} \rightarrow X$ be the tautological morphism.

Note that π admits a section $\mathcal{E}$, which is contracted by φ to the point x . Consider the blowing-up $\sigma : X' \rightarrow X$ of X at x . Via [IN, Lemma 4.3], the universal property of the blowing-up shows that we have a morphism $\psi : \mathcal{F} \rightarrow X'$, such that $\sigma \circ \psi = \varphi$. So we have the following diagram

$$\begin{array}{ccc} \mathcal{F} & \xrightarrow{\psi} & X' \\ \pi \downarrow & \searrow \varphi & \downarrow \sigma \\ \mathcal{C} & & X. \end{array}$$

In particular, ψ maps the section $\mathcal{E}$ to E , the exceptional locus of σ . Let $\psi_0 : \mathcal{E} \rightarrow E$ be the restriction of ψ .

Step II. ψ_0 is generically finite and surjective; in particular, $\dim(\mathcal{C}) = n - 1$.

This follows from an elementary, but very useful remark, due to Kebekus [Ke1], proof of Theorem 3.4. Indeed, consider a general point $p \in E \simeq \mathbb{P}(T_x^*(X))$ and let

$C \subset \mathcal{E} \simeq \mathcal{C}$ be a curve such that $\psi_0(C) = p$. Take $\tilde{C}$, the normalization of C , and consider the surface S over $\tilde{C}$ which is got by base-change from our family. Let $D \subset S$ be the induced section for $\pi_S : S \rightarrow \tilde{C}$ and note that S is smooth along D by our hypothesis. The restriction of the tangent morphism of $\varphi_S : S \rightarrow X$ induces a morphism $T_{\varphi_S} : N_{D|S} \rightarrow l_p \simeq \mathbb{C}$, where $N_{D|S}$ is the (geometric) normal bundle of D in S and l_p is the line in $T_x(X)$ corresponding to $p \in E$. Since $N_{D|S}$ is not trivial, the above map has a zero. The corresponding curve of the family is singular at x . This is a contradiction. To conclude, note also that $\dim(\mathcal{F}) \geq n$, so $\dim(\mathcal{E}) \geq n - 1$, while $\dim(E) = n - 1$. In particular, $\dim(\mathcal{C}) = n - 1$ and φ (or ψ) is generically finite.

Step III. Conclusion.

Note first that the infinitesimal uniqueness property of our family translates into saying that the map ψ_0 is birational. Observe also that we have $\psi^{-1}(E) = \mathcal{E}$. Now, use properness of ψ and the fact that ψ_0 is birational to deduce that ψ is birational. We have already seen that $\mathcal{C}$ is birational to $E \simeq \mathbb{P}^{n-1}$. Replacing $\mathcal{F}$ by its normalization $\tilde{\mathcal{F}}$ and using the section $\mathcal{E}$, we find that $\tilde{\mathcal{F}}$ is generically a $\mathbb{P}^1$ -bundle over $\mathcal{C}$. Thus X is rational, being birational to $\tilde{\mathcal{F}}$. $\square$

The next result is a “more global” version of Theorem 1.3; it improves [IN, Theorem 4.2]. In the terminology of [IV], (ii) says that the model (X, C) is equivalent to $(\mathbb{P}^n, \text{line})$. Let us recall the following definition from [BBI]:

Definition 1.4 ([BBI]). A smooth rational curve $C \subset X$, where X is a projective manifold of dimension n , is a *quasi-line* if $N_{C|X} \simeq \bigoplus_1^{n-1} \mathcal{O}_{\mathbb{P}^1}(1)$.

Theorem 1.5. *Let X be a projective manifold and let $\mathcal{C}$ be a smooth maximal x -covering family of it. Let $\mathcal{D} \subset \mathcal{C}$ denote the family of reducible cycles in $\mathcal{C}$.*

- (i) *If the general member of $\mathcal{C}$ is smooth, it is a quasi-line (this is always the case if $\dim(X) \geq 3$).*
- (ii) *Assume moreover that the general member of $\mathcal{C}$ meets the locus of $\mathcal{D}$ only at x (this is true, e.g. if $\text{codim}_{\mathcal{C}}(\mathcal{D}) \geq 2$). Then, for any quasi-line $C \in \mathcal{C}$, there are a neighbourhood U of C in X and a neighbourhood V of a line $l \subset \mathbb{P}^n$, together with an isomorphism $U \simeq V$, sending C to l .*

Proof. We use the setting and notation from the proof of Theorem 1.3.

(i) Let $C \in \mathcal{C}$ be a general smooth curve. As $\mathcal{C}$ is a maximal x -covering family, $N_{C|X}$ is ample. In Step II from the proof of Theorem 1.3 we saw that $\dim(\mathcal{C}) = n - 1$, so C is a quasi-line.

(ii) We first prove the following:

Claim: There is a unique quasi-line from our given family passing through a general point of X (i.e. the map φ is birational).

Let $D \subset X'$ be the locus of points belonging to the images via ψ of reducible cycles from $\mathcal{C}$. By (i), a general curve C from our family $\mathcal{C}$ is a quasi-line in X . By the assumption in (ii), if $\tilde{C}$ denotes its proper transform on X' , we have $\tilde{C} \cap D = \emptyset$ for C general. So any point $c \in \pi(\psi^{-1}(\tilde{C}))$ represents an irreducible member of

the family $\mathcal{C}$. Now, the birationality of the map φ follows from the proof of [Ko, V.3.7.5.], a result due to Miyaoka. So the Claim is proved.

The Claim allows us to apply [Io2, Theorem 2.5] in order to get the conclusion in (ii). Finally, observe that $\text{codim}_{\mathcal{C}}(\mathcal{D}) \geq 2$ implies $\text{codim}_{X'}(D) \geq 2$. Moreover, if C is a general member of $\mathcal{C}$, the normal bundle of $\tilde{C}$ in X' is trivial; see, for instance, [IN, Lemma 1.6]. So, by [Ko, II.3.7], we have $\tilde{C} \cap D = \emptyset$ for C general. $\square$

2. CLASSIFICATION OF CONIC-CONNECTED MANIFOLDS

Definition 2.1 (cf. also [KS]). A smooth irreducible non-degenerate projective variety $X \subset \mathbb{P}^N$ is said to be a *conic-connected manifold*, briefly a *CC-manifold*, if through two general points of X there passes an irreducible conic contained in X .

The following classification theorem is our main result; as CC-manifolds are stable under isomorphic projection, we may assume X to be linearly normal.

Theorem 2.2. *Let $X \subset \mathbb{P}^N$ be a smooth irreducible linearly normal non-degenerate CC-manifold of dimension n . Then either $X \subset \mathbb{P}^N$ is a Fano manifold with $\text{Pic}(X) \simeq \mathbb{Z}\langle \mathcal{O}_X(1) \rangle$ and of index $i(X) \geq \frac{n+1}{2}$, or it is projectively equivalent to one of the following:*

- (i) $\nu_2(\mathbb{P}^n) \subset \mathbb{P}^{\frac{n(n+3)}{2}}$.
- (ii) *The projection of $\nu_2(\mathbb{P}^n)$ from the linear space $\langle \nu_2(\mathbb{P}^s) \rangle$, where $\mathbb{P}^s \subset \mathbb{P}^n$ is a linear subspace; equivalently $X \simeq \text{Bl}_{\mathbb{P}^s}(\mathbb{P}^n)$ embedded in $\mathbb{P}^N$ by the linear system of quadric hypersurfaces of $\mathbb{P}^n$ passing through $\mathbb{P}^s$; alternatively $X \simeq \mathbb{P}_{\mathbb{P}^r}(\mathcal{E})$ with $\mathcal{E} \simeq \mathcal{O}_{\mathbb{P}^r}(1)^{\oplus n-r} \oplus \mathcal{O}_{\mathbb{P}^r}(2)$, $r = 1, 2, \dots, n-1$, embedded by $|\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)|$. Here $N = \frac{n(n+3)}{2} - \binom{s+2}{2}$ and s is an integer such that $0 \leq s \leq n-2$.*
- (iii) *A hyperplane section of the Segre embedding $\mathbb{P}^a \times \mathbb{P}^b \subset \mathbb{P}^{N+1}$. Here $n \geq 3$ and $N = ab + a + b - 1$, where $a \geq 2$ and $b \geq 2$ are such that $a + b = n + 1$.*
- (iv) $\mathbb{P}^a \times \mathbb{P}^b \subset \mathbb{P}^{ab+a+b}$ *Segre embedded, where a, b are positive integers such that $a + b = n$.*

Corollary 2.3. *CC-manifolds are Fano and have second Betti number $b_2 \leq 2$; when $b_2 = 2$, they are also rational.*

Proof of Theorem 2.2. Fix a general point $x \in X$. By definition of CC-manifold, there exists a (maximal) x -covering family of conics on X ; we fix one such family, which we denote by $\mathcal{C}_x$.

Assume first that there exist lines l_1 and l_2 such that $l_1 \cup l_2$ belongs to $\mathcal{C}_x$.

Let $[C]$ denote the numerical class in $\text{Num}(X)$ of the curve C . Thus, if we let $[Q]$ be the class of a general conic from the family $\mathcal{C}_x$, we have $[Q] = [l_1] + [l_2]$. Let V_i , $i = 1, 2$, be the irreducible proper component of $\text{Chow}_{1,1}(X)$ to which l_i belongs, notation as in [Ko].

As in the first section, let us call the family V_i covering if its curves fill up X . Observe that, by the generality of the point $x \in X$, at least one of the families V_1, V_2 is so.

For each covering family V_i , define a relation R_i on X by saying that two points $x_1, x_2 \in X$ are equivalent if there exists a connected chain of lines from the family V_i joining the points x_1 and x_2 . This is the $\langle(U_i)_1\rangle$ -relation, $i = 1, 2$, for the normal form associated to the proper connected universal family $U_i \rightarrow V_i$ [Ko, Sections IV.4, IV.4.4 and IV.4.8.3].

Theorem IV.4.16 of [Ko] (see also [De, Theorem 5.9]) yields the existence of an open subset $X_i^0 \subseteq X$ and of a proper morphism with connected fibers $\pi_i : X_i^0 \rightarrow Z_i^0$ such that the $\langle(U_i)_1\rangle$ -relation restricts to an equivalence relation on X_i^0 and such that, for every $z \in Z_i^0$, $\pi_i^{-1}(z)$ coincides with the equivalence relation class defined by V_i , $i = 1, 2$. Since the family is covering the equivalence class of a point in X_i^0 does not reduce to the point itself. Therefore the equivalence class of a general point $x \in X$ is scheme theoretically a smooth irreducible positive dimensional projective algebraic variety $F_i^x = \pi_i^{-1}(\pi_i(x))$. Let $a_i = \deg N_{l_i|X}$. Since we are assuming V_i covering, $a_i \geq 0$; the locus of lines (from the family determined by l_i) passing through a general point x will be denoted by C_i^x . It is a cone of dimension $a_i + 1 \geq 1$. Clearly, we have $C_i^x \subseteq F_i^x$, so $\dim(F_i^x) \geq a_i + 1$ and equality holds if and only if F_i^x is a linear space.

Case I. Assume that $[l_1] = [l_2]$.

Let $[l] = [l_i]$, $i = 1, 2$, and let $Q \in \mathcal{C}_x$ be a general conic. Let R_x be the locus of points on X which can be joined to x by a connected chain of lines whose numerical class is $[l]$. If $R_x \subsetneq X$, reasoning as above, we could construct an open subset $X^0 \subseteq X$ and a proper morphism with connected fibers $\pi : X^0 \rightarrow Z^0$ such that, for every $z \in Z^0$, $\pi^{-1}(z)$ coincides with the equivalence relation class defined by the irreducible proper component of $\text{Chow}_{1,1}(X)$ to which l belongs. From this it would follow the existence of an effective divisor D passing through a general point $z \in X$ and such that $D \cdot l = 0$. This would imply $D \cdot Q = 2D \cdot l = 0$, which is clearly impossible. In conclusion $R_x = X$, so that the Picard number of X is one by [Ko, IV.3.13.3]. In this case, clearly $\text{Pic}(X) = \mathbb{Z}\langle\mathcal{O}(1)\rangle$. Moreover, Q has ample normal bundle, so $2i(X) = -K_X \cdot Q \geq n + 1$, giving $i(X) \geq \frac{n+1}{2}$.

From now on, assume that $[l_1] \neq [l_2]$, so that $V_1 \neq V_2$.

Case II. Assume that both V_1 and V_2 are covering. So $a_i \geq 0$, $i = 1, 2$.

Keeping the notation introduced above, let us observe that

$$\begin{aligned}
 \dim(F_1^x) + \dim(F_2^x) &\geq \dim(C_1^x) + \dim(C_2^x) = a_1 + 1 + a_2 + 1 \\
 (2.1) \quad &= -K_X \cdot l_1 - 1 + (-K_X) \cdot l_2 - 1 \\
 &= -K_X \cdot Q - 2 = \dim(\mathcal{C}_x) \geq n - 1.
 \end{aligned}$$

Moreover, since $[l_1] \neq [l_2]$, an R_1 -equivalence class and an R_2 -equivalence class cannot intersect along a curve, see [Ko, IV.3.13.3]. Note that we shall use this systematically in what follows.

Assume first that $\dim(F_1^x) + \dim(F_2^x) = n$. If $y \in X$ is another general point, using the rational maps π_i and the fact that $\dim(F_1^x \cap F_2^x) \leq 0$, we see that $F_1^x \cap F_2^y \neq \emptyset$. In particular, any point of X may be joined to x by a connected chain of rational curves from the families V_1, V_2 . It follows from [Ko, IV.3.13.3]

that $N_1(X)_\mathbb{Q} = \langle [l_1], [l_2] \rangle$. Moreover, by [BCD, Proposition 1], both rays $\mathbb{R}^+[l_i]$ are extremal since R_i -equivalence classes are equidimensional, for $i = 1, 2$.

Assume next that $\dim(F_1^x) + \dim(F_2^x) = n - 1$. By (2.1) this gives $a_1 + a_2 + 2 = n - 1$. Denote by E_i^x the (closed) locus of points lying on rational curves from the family V_j meeting F_i^x , for $i \neq j$. We have $\dim(E_i^x) = a_1 + a_2 + 2 = n - 1$, $i = 1, 2$. It follows from the definition of E_i^x that $E_i^x \cdot l_j = 0$ for $i \neq j$. We must have $E_i^x \cdot l_i > 0$; otherwise we get $E_i^x \cdot Q = 0$, which is absurd since E_i^x is effective and x is general. From this it follows again that any point of X may be joined to x by a connected chain of rational curves from the families V_1, V_2 , so that $N_1(X)_\mathbb{Q} = \langle [l_1], [l_2] \rangle$. We claim that both rays $\mathbb{R}^+[l_1], \mathbb{R}^+[l_2]$ are extremal. Take a curve $C \subset X$ and write its class in $N_1(X)_\mathbb{Q}$ as $[C] = a_1[l_1] + a_2[l_2]$, $a_1, a_2 \in \mathbb{Q}$. We may assume (up to a permutation of indices) that $a_1 \geq 0$. If $E_2^x \cdot C \geq 0$, it follows $a_2 \geq 0$ and we are done. If not, we must have $C \subset E_2^x$. But in this case, a variant of “bend and break” shows that we may write $[C] = \alpha_1[l_1] + \alpha_2[l_2]$, with $\alpha_2 \geq 0$. To the best of our knowledge, this last trick was first noticed in [BSW, Lemma 1.4.5].

In conclusion, the cone of effective 1-cycles $NE(X) = \overline{NE(X)} = \mathbb{R}^+[l_1] + \mathbb{R}^+[l_2]$ and $-K_X$ is ample by the Kleiman criterion. Thus X is a Fano manifold with $\text{Pic}(X) \simeq \mathbb{Z} \oplus \mathbb{Z}$.

There exist two contractions $f_i : X \rightarrow Z_i$ of the extremal rays $\mathbb{R}^+[l_i]$, where Z_i is a normal projective variety with $\text{Pic}(Z_i) \simeq \mathbb{Z}$ for $i = 1, 2$. Write, for simplicity, $F_i = F_i^x$ and $C_i = C_i^x$ for $i = 1, 2$.

From (2.1) it follows that exactly one of the following subcases (a), (b) or (c) may arise.

- (a) $F_i = C_i \simeq \mathbb{P}^{a_i+1}$, $i = 1, 2$, $\dim(F_1) + \dim(F_2) = n$, so $a_1 + a_2 + 2 = n$,
- (b) $F_i = C_i \simeq \mathbb{P}^{a_i+1}$, $i = 1, 2$, $\dim(F_1) + \dim(F_2) = n - 1$, so $a_1 + a_2 + 2 = n - 1$,
- (c) $F_1 = C_1 \simeq \mathbb{P}^{a_1+1}$, $C_2 \subsetneq F_2$, $\dim(F_1) + \dim(F_2) = n$, so $a_1 + a_2 + 3 = n$.

Subcase (a). The linear spaces F_1, F_2 intersect transversally in a point. It follows that the map $f_1 \times f_2 : X \rightarrow Z_1 \times Z_2 \simeq \mathbb{P}^{a_2+1} \times \mathbb{P}^{a_1+1}$ is finite and birational, hence an isomorphism. This leads to case (iv).

Subcase (b). At least one of the contractions f_1, f_2 is equidimensional. Otherwise we may find two fibres F_1, F_2 such that $\dim(F_1 \cap F_2) \geq 1$. Assume that f_1 is equidimensional. From [Fu1, Lemma 2.12] or [Io1, p. 467], it follows that $f_1 : X \rightarrow Z_1$ is a $\mathbb{P}^{a_1+1}$ -bundle. In particular, Z_1 is smooth. At this point, we want to apply the characterization of projective space due to Cho–Miyaoka–Shepherd-Barron [CMSB] to see that $Z_1 \simeq \mathbb{P}^{a_2+2}$. As the proof of this important criterion is quite involved, we would like to explain how, due to our particular context, the result follows from the simpler argument in [Ke2]. Note that smoothness of Z_1 is important for this simplification. Let Q be a conic from our family $\mathcal{C}_x$. If $Q = l_1 \cup l_2$ is reducible, we have $f_1(Q) = f_1(l_2)$ and the restriction $f_1|_{l_2} : l_2 \rightarrow f_1(l_2)$ is an isomorphism. Let now Q be irreducible and let Π denote the plane it spans. We claim that $f_1|_Q : Q \rightarrow f_1(Q)$ is birational. This is seen by looking at the intersection of the linear spaces Π and a fibre F_1 . If $x \in X$ is a very general point,

consider the family of irreducible reduced rational curves $f_{1*}(Q)$, where $Q \in \mathcal{C}_x$. The argument in [Ke2] applies to this family, yielding that $Z_1 \simeq \mathbb{P}^{a_2+2}$ and $f_1(Q)$ is a line.

A general conic Q from our family has ample normal bundle and $\deg N_{Q|X} = a_1 + a_2 + 2 = n - 1$, so Q is a quasi-line. Now fix a general conic Q_0 and let $L := f_1(Q_0)$, a line in $Z_1 = \mathbb{P}^{a_2+2}$. Let $T := f_1^{-1}(L)$. Fix also two general points $t, t' \in T$. Any conic Q passing through t, t' is contained in T , because $f_1(Q)$ is a line. The standard exact sequence of normal bundles

$$0 \rightarrow N_{Q|T} \rightarrow N_{Q|X} \rightarrow N_{T|X|Q} \rightarrow 0$$

shows that Q is a quasi-line in T . Therefore, we may apply [IV, Proposition 4.1] to deduce that (up to a Stein factorization) the only non-constant non-finite morphisms defined on T are $f_1|_T$ and $\sigma : T \rightarrow \mathbb{P}^{a_1+2}$, a blowing-up sending Q to a line of $\mathbb{P}^{a_1+2}$. Consider now the restriction $r := f_2|_T : T \rightarrow Z_2$. As $\dim(T) = a_1 + 2$, by the above considerations, r is surjective. We claim that r is birational. Otherwise, take two distinct points t_1, t_2 in a general fibre, $F_2 \cap T$, of r . The line $\langle t_1, t_2 \rangle \subset F_2$ meets T in two points, so it is contained in T by the above reasoning. This is absurd since r is generically finite. Moreover, r is not an isomorphism, as $\text{Pic}(Z_2) \simeq \mathbb{Z}$ and $\text{Pic}(T) \simeq \mathbb{Z} \oplus \mathbb{Z}$. Therefore, $r = \sigma$, in particular $Z_2 \simeq \mathbb{P}^{a_1+2}$ and $f_2(Q)$ is a line.

To conclude subcase (b), we write the hyperplane section of X as $H = H_1 + H_2$, with $H_i \cdot l_j = 1 - \delta_{ij}$. We have seen that $f_i^*(D_i) = H_i$, where D_i is the hyperplane divisor of Z_i for $i = 1, 2$. The map $f := f_1 \times f_2 : X \rightarrow \mathbb{P}^{a_2+2} \times \mathbb{P}^{a_1+2}$ is given by a linear system $\Lambda \subseteq |H|$. Let $X' := f(X)$, which is a divisor on $Z := \mathbb{P}^{a_2+2} \times \mathbb{P}^{a_1+2}$, and note that $f : X \rightarrow X'$ is finite and birational by construction. Since $H \cdot l_i = 1$ for $i = 1, 2$, the general fiber of f_i is mapped by f onto a hyperplane of Z_j , for $i \neq j$. Therefore, the divisor X' has type $(1, 1)$ in $\text{Pic}(Z)$ and f induces an isomorphism on each fiber of f_1 . Thus $f : X \rightarrow X'$ is an isomorphism and we get case (iii) of the theorem.

Subcase (c). In this case, both f_1 and f_2 are equidimensional and $\dim(Z_2) = n - a_2 - 2 = a_1 + 1 = \dim(F_1)$. Using the notation and the reasoning above we get $\dim(T) = a_1 + 2$, so $f_2|_T : T \rightarrow Z_2$ cannot be birational. This shows subcase (c) does not happen.

Case III. Assume that the family V_1 is covering and V_2 is not.

Now we can also assume that the maximal x -covering family $\mathcal{C}_x$ is smooth (otherwise we are in Case II for some different covering family V_2), so we may use Theorem 1.5. In particular, the general conic in $\mathcal{C}_x$ is a quasi-line.

We consider separately the subcases:

III₁. C_1^x has codimension 1 in X .

III₂. C_1^x has codimension ≥ 2 in X .

Case III₁. Reasoning as in Case II, we deduce that either $X = F_1^x \supsetneq C_1^x$, or $F_1^x = C_1^x$ is a linear subspace in $\mathbb{P}^N$. In the first case, $\text{Pic}(X) = \mathbb{Z}\langle \mathcal{O}(1) \rangle$ and we would fall in Case I, contrary to our assumption. In the second case, F_1^x has trivial normal bundle, so that $X \subset \mathbb{P}^N$ is easily seen to be a scroll over $\mathbb{P}^1$, say

$\pi : X \rightarrow \mathbb{P}^1$, having F_1^x as general fiber. We may write $X \simeq \mathbb{P}(\mathcal{E})$, with $\mathcal{E} = \pi_*(\mathcal{O}_X(1))$ a rank- n ample vector bundle on $\mathbb{P}^1$. Since the general conic $Q \in \mathcal{C}_x$ is a quasi-line, $-K_X \cdot Q = n + 1$. Using the formula for the canonical class of $\mathbb{P}(\mathcal{E})$ and the relation $F_1^x \cdot Q = 1$, we obtain $\deg(\det \mathcal{E}) = n + 1$. As $\mathcal{E}$ is ample, its splitting-type must be $(1, \dots, 1, 2)$ and we get case (ii) with $r = 1$ (or $s = n - 2$).

Case III₂.

Claim: If Q is a general member of $\mathcal{C}_x$, Q meets the locus of degenerate conics only at x .

Let $D \subset X$ denote the locus of V_2 . By [Ko, II.3.7], to prove the Claim it is sufficient to show that $D \cap Q = \emptyset$ when D is a prime divisor. In this case C_1^x is also irreducible and the locus of degenerate conics is $C_1^x \cup D$.

The strategy for proving the above Claim is in two steps. First, as in Case II, we show that the rationally connected fibration determined by the covering family V_1 is a $\mathbb{P}^{a_1+1}$ -bundle over $\mathbb{P}^{n-a_1-1}$. Next, we deduce from this that a general conic through x does not intersect D , proving the Claim. We now proceed with the details.

Let y be a general point of $F_1^x \cap D$. As before, let C_2^y be the cone of lines from V_2 passing through y . The geometrical description of D yields

$$\dim(C_1^x \cap D) + \dim(C_2^y) = \dim(D) = n - 1.$$

Since

$$\dim(C_1^x \cap D) = \dim(C_1^x) - 1 = -K_X \cdot l_1 - 2,$$

we find

$$\dim(C_2^y) = n - 1 + K_X \cdot l_1 + 2 = -K_X \cdot Q + K_X \cdot l_1 = -K_X \cdot l_2 \quad \text{for } y \in D \text{ general.}$$

It follows

$$(2.2) \quad \dim(C_2^y) \geq -K_X \cdot l_2 \quad \text{for any } y \in D.$$

By generality of x and y we may assume that $(F_1^x \cap D) \cap C_2^y$ meets the smooth locus of D . We deduce

$$\begin{aligned} 0 &\geq \dim((F_1^x \cap D) \cap C_2^y) \geq \dim(F_1^x \cap D) + \dim(C_2^y) - (n - 1) \\ &\geq \dim(C_1^x \cap D) + \dim(C_2^y) - (n - 1) = 0. \end{aligned}$$

In particular, $\dim(F_1^x \cap D) = \dim(C_1^x \cap D)$. Thus, $F_1^x = C_1^x = \mathbb{P}^{a_1+1}$, $a_1 = \deg(N_{l_1|X}) \geq 0$. Next we prove that the rationally connected fibration determined by V_1 is a $\mathbb{P}^{a_1+1}$ -bundle over some manifold Z . Indeed, let $\pi : X^0 \rightarrow Z^0$ be our fibration, which is generically a $\mathbb{P}^{a_1+1}$ -bundle by the above. We show that $X^0 = X$ and π is equidimensional. Using the notation in [BCD, Proposition 1], assume that $B \neq \emptyset$. We claim that $B \cap D \neq \emptyset$. Indeed, if $b \in B \setminus D$, by definition of B there exists a line $l_1 \in V_1$ contained in B and containing b . Since $D \cdot l_1 > 0$, the last claim follows. By [BCD, Proposition 1], there is some equivalence class $\overline{F_1}$ contained in B , passing through a point $b \in B \cap D$ and having $\dim(\overline{F_1}) > a_1 + 1$. We deduce, using (2.2)

$$\dim(\overline{F_1} \cap C_2^b) \geq \dim(\overline{F_1}) + \dim(C_2^b) - n > a_1 + 1 - K_X \cdot l_2 - n = 0.$$

This would imply $[l_1] = [l_2]$, a contradiction. So, $B = \emptyset$ and π is equidimensional. As in Case II, subcase (b), it follows that $\pi : X \rightarrow Z$ is a $\mathbb{P}^{a_1+1}$ -bundle and $Z \simeq \mathbb{P}^{n-a_1-1}$. Moreover, a general conic through x is sent isomorphically by π onto a line in Z .

Let us fix Q , a general conic through x . Set $L := \pi(Q) \subset \mathbb{P}^{n-a_1-1}$ and let $T := \pi^{-1}(L) \rightarrow L$ be the associated $\mathbb{P}^{a_1+1}$ -bundle over $\mathbb{P}^1$. The conic Q is contained in T . Reasoning as in Case II, subcase (b), it is immediate to see that T is a CC-manifold, belonging to the class described in Case III₁ above. In particular there is a map $\sigma : T \rightarrow \mathbb{P}^{a_1+2}$ which is a blowing-up along a linear $\mathbb{P}^{a_1}$. General conics in T are identified via σ to lines in $\mathbb{P}^{a_1+2}$ and $D \cap T$ is the exceptional divisor of σ . Thus $Q \cap D = Q \cap (D \cap T) = \emptyset$ and the Claim is finally proved.

By Theorem 1.5(ii), there exists a birational map $\varphi : \mathbb{P}^n \dashrightarrow X \subset \mathbb{P}^N$ given by a linear system $\Lambda \subseteq |\mathcal{O}_{\mathbb{P}^n}(2)|$, inducing an isomorphism between a neighbourhood of a line in $\mathbb{P}^n$ and a neighbourhood of a conic $Q \in \mathcal{C}_x$. In particular φ is an isomorphism outside the base locus of Λ , which we denote by Y . We claim that, if $Y \neq \emptyset$, Y is a linear space $\mathbb{P}^s$ with $0 \leq s \leq n-3$.

Suppose there exist two distinct points $y_1, y_2 \in Y$. The linear system of quadric hypersurfaces Λ cannot separate points along the line $\langle y_1, y_2 \rangle$. Therefore $\langle y_1, y_2 \rangle \subseteq Y$. The infinitesimal version of the same argument shows Y is reduced, so it is a linear space. Thus we are in one of the cases described in (ii).

Finally, if all conics in $\mathcal{C}_x$ are irreducible, we get case (i) of the theorem. This follows directly from Theorem 1.5(ii). $\square$

Conversely, from the recent paper [BH], we see that the condition $i \geq \frac{n+1}{2}$ in Theorem 2.2 is optimal.

Proposition 2.4. *Let $X \subset \mathbb{P}^{n+r}$ be a smooth non-degenerate complete intersection of dimension n and multi-degree $(d_1, d_2, \dots, d_r)$. If we assume that $n \geq 2 \sum_1^r d_i - 2r - 1$, then X is a CC-manifold.*

Proof. Note that the inequality in the statement means that X is Fano, of index at least $\frac{n+1}{2}$. Replacing X by a general hyperplane section, we may assume that equality holds. Therefore, we may apply [BH, Corollary 1.2] to conclude. $\square$

The next result, essentially due to Hwang–Kebekus [HK, Theorem 3.14], shows that any Fano manifold of high index is conic-connected. Note that we slightly improve the bound on the index given in [HK]. The first part of the proposition is well known.

Proposition 2.5 (cf. [HK]). *Let $X \subset \mathbb{P}^N$ be a Fano manifold with $\text{Pic}(X) \simeq \mathbb{Z}\langle H \rangle$ and $-K_X = i(X)H$, H being the hyperplane section and $i(X)$ the index of X .*

- (i) *If $i(X) > \frac{n+1}{2}$, then $X \subset \mathbb{P}^N$ is ruled by lines and for general $x \in X$ the Hilbert scheme of lines through x , $Y_x \subset \mathbb{P}(T_x^*(X)) = \mathbb{P}^{n-1}$, is smooth. If $i(X) \geq \frac{n+3}{2}$, Y_x is also irreducible. Let SY_x be the secant variety of Y_x in $\mathbb{P}^{n-1}$.*
- (ii) *If $i(X) \geq \frac{n+3}{2}$ and $SY_x = \mathbb{P}^{n-1}$, then $X \subset \mathbb{P}^N$ is a CC-manifold.*

(iii) If $i(X) > \frac{2n}{3}$, then $X \subset \mathbb{P}^N$ is a CC-manifold.

Proof. By a theorem of Mori [Mo1] the variety $X \subset \mathbb{P}^N$ is ruled by a family of rational curves $\mathcal{L}$ such that for $L \in \mathcal{L}$ we have $-K_X \cdot L \leq n + 1$. It follows that $X \subset \mathbb{P}^N$ is ruled by lines, that is $H \cdot L = 1$. As is well known, see [Ko, II.3.11.5] for a more general result, the Hilbert scheme of lines passing through x is smooth equidimensional and may be identified with a subscheme $Y_x \subset \mathbb{P}(T_x^*(X)) = \mathbb{P}^{n-1}$ of dimension $i(X) - 2$. Thus, if $i(X) - 2 \geq \frac{n-1}{2}$, $Y_x \subset \mathbb{P}^{n-1}$ is irreducible.

Part (ii) follows from [HK, Theorem 3.14].

To prove part (iii), first observe that $i(X) \geq \frac{n+3}{2}$, unless $n \leq 6$. If $n \leq 6$, $i(X) > \frac{2n}{3}$ gives $i(X) \geq n - 1$, so the conclusion follows by the classification of del Pezzo manifolds (see [Fu2]). To conclude, using part (ii), it is enough to see that $SY_x = \mathbb{P}^{n-1}$.

By [Hw, Theorem 2.5], the variety $Y_x \subset \mathbb{P}^{n-1}$ is non-degenerate and by hypothesis $n - 1 < \frac{3i(X)-6}{2} + 2 = \frac{3\dim(Y_x)}{2} + 2$, so that $SY_x = \mathbb{P}^{n-1}$ by Zak Linear Normality Theorem; see [Za, V.1.13]. $\square$

ACKNOWLEDGEMENTS

Both authors are grateful to the organizers of the Conference “Projective varieties with unexpected properties”, that took place in Siena, between 8–13 of June 2004. Our collaboration started while taking part in this very pleasant and fruitful mathematical event.

We thank Tommaso de Fernex and Ciro Ciliberto for a critical reading of previous versions of our paper.

REFERENCES

- [BBI] L. BĂDESCU, M.C. BELTRAMETTI, P. IONESCU, Almost-lines and quasi-lines on projective manifolds, in *Complex Analysis and Algebraic Geometry* (in memory of Michael Schneider), de Gruyter, 2000, pp. 1–27.
- [BSW] M.C. BELTRAMETTI, A.J. SOMMESE, J. WIŚNIEWSKI, Results on varieties with many lines and their applications to adjunction theory, in *Complex Algebraic Varieties (Bayreuth, 1990)*, Lecture Notes in Math., 1507, Springer, 1992, pp. 16–38.
- [BCD] L. BONAVERO, C. CASAGRANDE, S. DRUEL, On covering and quasi-unsplit families of curves, *J. Eur. Math. Soc.* **9** (2007), 45–57.
- [BH] L. BONAVERO, A. HÖRING, Counting conics in complete intersections, Preprint (2008), math. AG/0804162.
- [Ca] F. CAMPANA, Coréduction algébrique d’un espace analytique faiblement kählerien compact, *Invent. Math.* **63** (1981), 187–223.
- [CMSB] K. CHO, Y. MIYAOKA, N. SHEPHERD-BARRON, Characterizations of projective space and applications to complex symplectic manifolds, in *Higher Dimensional Birational Geometry (Kyoto, 1997)*, Adv. Stud. Pure Math., vol. 35, Math. Soc. Japan, 2002, pp. 1–88.
- [De] O. DEBARRE, *Higher-Dimensional Algebraic Geometry*, Universitext, Springer, 2001.
- [Fu1] T. FUJITA, On polarized manifolds whose adjoint bundles are not semipositive, in *Algebraic Geometry (Sendai, 1985)*, Adv. Stud. Pure Math., vol. 10, North-Holland, 1987, pp. 167–178.
- [Fu2] T. FUJITA, *Classification Theories of Polarized Varieties*, London Math. Soc. Lecture Note Ser., vol. 155, Cambridge Univ. Press, 1990.

- [Hw] J.M. HWANG, Geometry of minimal rational curves on Fano manifolds, in *School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000)*, ICTP Lect. Notes, vol. 6, Abdus Salam Int. Cent. Theoret. Phys., 2001, pp. 335–393.
- [HK] J.M. HWANG, S. KEBEKUS, Geometry of chains of minimal rational curves, *J. Reine Angew. Math.* **584** (2005), 173–194.
- [Io1] P. IONESCU, Generalized adjunction and applications, *Math. Proc. Cambridge Philos. Soc.* **99** (1986), 457–472.
- [Io2] P. IONESCU, Birational geometry of rationally connected manifolds via quasi-lines, in *Projective Varieties with Unexpected Properties (Siena, 2004)*, de Gruyter, 2005, pp. 317–335.
- [IN] P. IONESCU, D. NAIE, Rationality properties of manifolds containing quasi-lines, *Internat. J. Math.* **14** (2003), 1053–1080.
- [IR] P. IONESCU, F. RUSSO, Varieties with quadratic entry locus. II, *Compositio Math.* **144** (2008), 949–962.
- [IV] P. IONESCU, C. VOICA, Models of rationally connected manifolds, *J. Math. Soc. Japan* **55** (2003), 143–164.
- [Ke1] S. KEBEKUS, Families of singular rational curves, *J. Algebraic Geom.* **11** (2002), 245–256.
- [Ke2] S. KEBEKUS, Characterizing the projective space after Cho, Miyaoka and Shepherd-Barron, in *Complex Geometry (Göttingen, 2000)*, Springer, 2002, pp. 147–155.
- [Ko] J. KOLLÁR, *Rational Curves on Algebraic Varieties*, Ergeb. Math. Grenzgeb. (3), vol. 32, Springer, 1996.
- [KMM] J. KOLLÁR, Y. MIYAOKA, S. MORI, Rationally connected varieties, *J. Algebraic Geom.* **1** (1992), 429–448.
- [KS] Y. KACHI, E. SATO, *Segre’s Reflexivity and an Inductive Characterization of Hyperquadrics*, Mem. Amer. Math. Soc., vol. 160, no. 763, 2002.
- [Mo1] S. MORI, Projective manifolds with ample tangent bundle, *Ann. of Math.* **110** (1979), 593–606.
- [Mo2] S. MORI, Threefolds whose canonical bundles are not numerically effective, *Ann. of Math.* **116** (1982), 133–176.
- [Ru] F. RUSSO, Varieties with quadratic entry locus. I, Preprint (2006), math. AG/0701889, to appear in Math. Ann.
- [Sc1] G. SCORZA, Sulla determinazione delle varietà a tre dimensioni di S_r ($r \geq 7$) i cui S_3 tangenti si tagliano a due a due, *Rend. Circ. Mat. Palermo* **25** (1908), 193–204; contained in *Opere Scelte*, vol. I, Ed. Cremonese, 1960, pp. 167–182.
- [Sc2] G. SCORZA, Sulle varietà a quattro dimensioni di S_r ($r \geq 9$) i cui S_4 tangenti si tagliano a due a due, *Rend. Circ. Mat. Palermo* **27** (1909), 148–178; contained in *Opere Scelte*, vol. I, Ed. Cremonese, 1960, pp. 252–292.
- [Za] F.L. ZAK, *Tangents and Secants of Algebraic Varieties*, Transl. Math. Monogr., vol. 127, Amer. Math. Soc., 1993.

UNIVERSITY OF BUCHAREST, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, 14 ACADEMIEI STR., RO-010014 BUCHAREST, AND INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O. BOX 1-764, RO-014700 BUCHAREST, ROMANIA

E-mail address: Paltin.Ionescu@imar.ro

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DEGLI STUDI DI CATANIA, VIALE A. DORIA, 6, 95125 CATANIA, ITALY

E-mail address: frusso@dmf.unict.it